

ACETAMINOPHEN DOSING (FOR EXAMPLE, TYLENOL®)

NOT recommended for babies less than 2 months old. Give every 4-6 hours as needed for fever or pain.

NOTE: Certain concentrations (mg/mL) of acetaminophen are being phased out. Please be sure to check the concentration on the bottle of acetaminophen before dosing.

Weight (pounds)	Weight (kilogram)	Children's Liquid (160mg/5mL)	Children's Dissolve Tabs (80mg tabs)
12-17 lbs	5.5-7.9 kg	2.5 mL (½ teaspoon)	---
18-23 lbs	8-10.9 kg	3.7 mL (¾ teaspoon)	---
24-35 lbs	11-15.9 kg	5 mL (1 teaspoon)	2 tabs (160 mg)
36-47 lbs	16-20.9 kg	7.5 mL (1½ teaspoons)	3 tabs (240 mg)
48-59 lbs	21-26.5 kg	10 mL (2 teaspoons)	4 tabs (320 mg)
60-71 lbs	26.6-32.2 kg	12.5 mL (2½ teaspoons)	5 tabs (400 mg)

IBUPROFEN DOSING (FOR EXAMPLE, MOTRIN® AND ADVIL®)

For infants and children OLDER than 6 months of age. Give every 6-8 hours as needed for fever or pain.

Weight (pounds)	Weight (kilogram)	Children's Liquid (100mg/5mL)	Junior Tabs (100mg)	Adult Tabs (200mg)
12-21 lbs	5.5-7.9 kg	2.5 mL (½ teaspoon)	---	---
22-33 lbs	10-14.9 kg	5 mL (1 teaspoon)	1 tab (100mg)	---
34-43 lbs	15-19.9 kg	7.5 mL (1½ teaspoons)	1 tab (100mg)	---
44-45 lbs	20-24.9 kg	10 mL (2 teaspoons)	2 tabs (200 mg)	1 tab (200 mg)
56-66 lbs	25-29.9 kg	12.5 mL (2½ teaspoons)	2 tabs (200 mg)	1 tab (200 mg)
67-88 lbs	30-39.9 kg	15 mL (3 teaspoons)	3 tabs (300mg)	---
89+ LBS	40+ kg	---	4 tabs (400 mg)	2 tabs (400 mg)

3248 e 9/14



DOSIS DE ACETAMINOFÉN (POR EJEMPLO, TYLENOL®)

NO se recomienda para bebés menores de 2 meses de edad. Dar medicamento cada 4-6 horas para aliviar síntomas de fiebre o dolor. **NOTA:** En algunas concentraciones (mg/ml) de acetaminofén las fases se separan. Asegúrese de verificar la concentración en el frasco de acetaminofén antes de administrarla.

Peso (libras)	Peso (kilos)	Jarabe líquido para niños (160mg/5mL)	Tabletas masticables (tableta de 80 mg)
12-17 libras	5.5-7.9 kilos	2.5 mL (½ cucharita)	---
18-23 libras	8-10.9 kilos	3.7 mL (¾ cucharita)	---
24-35 libras	11-15.9 kilos	5 mL (1 cucharita)	2 tabletas (160 mg)
36-47 libras	16-20.9 kilos	7.5 mL (1½ cucharita)	3 tabletas (240 mg)
48-59 libras	21-26.5 kilos	10 mL (2 cucharita)	4 tabletas (320 mg)
60-71 libras	26.6-32.2 kilos	12.5 mL (2½ cucharita)	5 tabletas (400 mg)

DOSIS DE IBUPROFENO (POR EJEMPLO, MOTRIN® Y ADVIL®)

Para niños mayores de 6 meses. Dar medicamento cada 6-8 horas para aliviar síntomas de fiebre o dolor.

Peso (libras)	Peso (kilos)	Jarabe líquido para niños (100mg/5 mL)	Tabletas para niños (100mg)	Pastillas para adultos (200mg)
12-21 libras	5.5-7.9 kilos	2.5 mL (½ cucharita)	---	---
22-33 libras	10-14.9 kilos	5 mL (1 cucharita)	1 tableta (100mg)	---
34-43 libras	15-19.9 kilos	7.5 mL (1½ cucharita)	1 tableta (100mg)	---
44-45 libras	20-24.9 kilos	10 mL (2 cucharita)	2 tableta (200 mg)	1 pastilla (200 mg)
56-66 libras	25-29.9 kilos	12.5 mL (2½ cucharita)	2 tableta (200 mg)	1 pastilla (200 mg)
67-88 libras	30-39.9 kilos	15 mL (3 cucharita)	3 tableta (300mg)	---
89+ libras	40+ kilos	---	4 tableta (400 mg)	2pastilla (400 mg)